

Supplementary material

Shared and connection-specific intrinsic interactions in the default mode network

Jessica Samogin ^{1*}, Quanying Liu^{1,2,3,4*}, Marco Marino⁵, Nicole Wenderoth^{1,2} and Dante Mantini^{1,5}

¹ *Research Center for Motor Control and Neuroplasticity, KU Leuven, 3001 Leuven, Belgium;*

² *Neural Control of Movement Laboratory, ETH Zurich, 8057 Zurich, Switzerland;*

³ *Division of Engineering and Applied Science, California Institute of Technology, Pasadena, United States;*

⁴ *Molecular Neurology Program, Huntington Medical Research Institutes, Pasadena, United States;*

⁵ *Brain Imaging and Neural Dynamics Research Group, IRCCS San Camillo Hospital, 30126 Venice, Italy.*

Supplementary Table 1 – Definition of seed areas used in the study. The anatomical name, its abbreviation and the coordinates in MNI space are provided for each seed.

<i>Seed name</i>	<i>Abbreviation</i>	<i>MNI coordinates</i>
Posterior cingulate cortex	PCC	[0, -66, 28]
Medial prefrontal cortex	MPFC	[0, 44, 10]
Left angular gyrus	LANG	[-57, -63, 17]
Right angular gyrus	RANG	[57, -63, 17]




Fig. S1 – hdEEG connectivity map for DMN for different frequency bands. Connectivity was calculated for LANG and RANG as seed regions (light blue dot), respectively, in delta (1-4 Hz), theta (4-8 Hz), alpha (8-13 Hz), beta (13-30 Hz) and gamma (30-80 Hz) bands. Spatial maps are shown in coronal, sagittal and axial sections. The significance level is set to $p < 0.05$, FDR corrected.